




**Modification to K-CEWS: A Doctor can adjust the acceptable parameter based on specific treatments or pre-existing medical conditions. Modifications must NEVER be used to normalise a clinically unstable patient**

Vital Sign	New accepted values and new K-CEWS score	Time and Date	Duration	Name and contact details
Reason:				
Reason:				
Reason:				

Blood Pressure (record and score when needed)				
	Date	Time		
Blood Pressure: Score systolic only	>130s	3		3
	120s	2		2
	110s	1		1
	100s	0		0
	90s			
	80s	1		1
	70s	3		3
	60s			
	50s			



Modified Glasgow Coma Score: Child				
Date				
Time				
Pupil Size	1	<b>Eye Response</b>		
		Open normally	4	
		React to speech/touch	3	
		React to pain	2	
	2	No response	1	
		<b>Verbal Response</b>		
	3	Alert, talking normally, like their usual ability	5	
		Less than usual words or irritable cry	4	
		Cries or makes vocal sounds to pain only	3	
	4	Moans or facial grimaces	2	
		None	1	
	5	<b>Best Motor Response</b>		
Spontaneous movement		6		
Localise stimuli		5		
Withdraws from pain		4		
Abnormal flexion		3		
Abnormal extension		2		
6	No response	1		
	<b>Total Score</b>			
Pupil Response: (R)Reactive (S)Sluggish (F)Fixed	Pupil Size	L		
	Pupil Response	L		
	Pupil Size	R		
	Pupil Response	R		
Any other signs of altered status (✓)	Nausea/vomiting			
	Headache			
	Photophobia			
	Double vision			

Respiratory Distress Assessment			
Observe	Mild	Moderate	Severe
Behaviour	Normal	Some intermittent irritability	Increasing irritability and/or tiring
Respiratory rate	Mildly increased (yellow '1' zone)	Increased (peach coloured '2' zone)	Marked increase (pink '3' zone) or a marked decrease
Use of accessory muscles	Nil to mild chest wall in-drawing	Some in-drawing seen: - Trachea - Below the sternum - Between the ribs - Below the ribs Nasal flaring	Severe in-drawing and effort to breathe noted in same places as Moderate. Marked nasal flaring Grunting heard when breathing out
Apnoeic episodes	None	May have a brief apnoea once	Increasing, frequent or prolonged
Feeding	Normal	Feeding poorly/less than normal	Unable to feed or latch
<b>FLACC Pain Scale</b>	<b>Score 0</b>	<b>Score 1</b>	<b>Score 2</b>
Face	Relaxed facial expression/smiles	Occasional grimace or frown	Frequent frown/clenched jaw
Legs	Normal position/relaxed	Uneasy/restless/tense	Rigid legs
Activity	Moves easily/lying quietly in normal position	Moving around bed a lot, moving arms, or legs or back and forth	Tight body, rigid arms or legs, lying very still
Cry	No cry	Moans/whimpers	Crying steadily
Consolability	Content/relaxed	Reassured by comfort measures	Difficult to comfort